arXiv:1108.2256v1 [math.FA] 10 Aug 2011

Functional Integral Representation for Relativistic Schrödinger Operator Coupled to a Scalar Bose Field with $P(\phi)$ Interaction

Toshimitsu Takaesu

Faculty of Mathematics, Kyushu University,
Fukuoka, 812-8581, Japan

Abstract. In this paper the system of a semi-relativistic particle interacting with a scalar Bose field is investigated. The ultraviolet cutoff condition is imposed on the Bose field. In the main theorem, the functional integral representation of the semi group generated by the total Hamiltonian with $P(\phi)$ interaction is obtained.

Key words : Relativistic Schrödinger operator, Quantum Field Theory, Spectral analysis, Lévy process, Gaussian random process.

1 Introduction

In this paper the system of a quantum particle interacting with a scalar Bose field is investigated. The partilcle's Hamiltonian is given the relativistic Schrödinger operator with potential

$$H_{\mathrm{p}} = \sqrt{-\triangle + M^2} - M + V \tag{1}$$

on the Hilbert space $L^2(\mathbf{R}^d_{\mathbf{x}})$, where $M > 0$ is the rest mass of the particle. For the stochastic analysis of H_{p}, the asymptotic behavior of the eigenvector is analyzed by the functional integral representation in [4], and the functional representation in electromagnetic potential is derived in [13, 15]. In this paper, we construct the functional integral representation according to [13]. For other results on the stochastic analysis of H_{p}, refer to e.g. [15, 16, 19].

A scalar Bose field is constructed by stochastic process which investigated in constructive quantum field theory (refer to e.g. [5, 24]). The field operators $\{\phi(f)\}_{f \in \mathcal{K}_{\mathrm{b}}}$ are defined by the Gaussan random process indexed by a Hilbert space $\mathcal{K}_{\mathrm{b}}$ on a probability space $(Q_{\mathrm{b}}, \mathfrak{B}_{\mathrm{b}}, P_{\mathrm{b}})$. The state space is given by $L^2(Q_{\mathcal{K}_{\mathrm{b}}})$ and the free Bose Hamiltonian H_{b} is defined by the differential second quantization of $\omega_{\mathrm{b}}(-i\nabla)$ where ω_{b} is non-negative and continuous function. Physically $\omega_{\mathrm{b}}(\mathbf{k}) \geq 0$ denotes the one-particle energy of the field with momentum $\mathbf{k}$. Thus the triplet $\left(L^2(Q_{\mathcal{K}_{\mathrm{b}}}), H_{\mathrm{b}}, \{\phi(f)\}_{f \in \mathcal{K}_{\mathrm{b}}}\right)$ of the scalar Bose field is defined.

The system of semi-relativistic particles coupled to a scalar Bose field is defined as follows. The state space is given by $\mathcal{H} = L^2(\mathbf{R}^d_{\mathbf{x}}) \otimes L^2(Q_{\mathrm{b}}) \simeq \int^{\oplus}_{\mathbf{R}^d} L^2(Q_{\mathrm{b}}) d\mathbf{x}$ where $\int^{\oplus}$ denotes the fibre direct integral. The free Hamiltonian is defined by $H_0 = H_\mathrm{p} \otimes I + I \otimes H_\mathrm{b}$ and the total Hamiltonian by

$$H_\kappa \;=\; H_0 \dotplus \kappa \int_{\mathbf{R}^d}^{\oplus} P(\phi(\rho_\mathbf{x}))d\mathbf{x} \tag{2}$$

where $\dotplus$ denotes the form sum, $P(\lambda) = \sum_{j=1}^{2n} c_j \lambda^j$, $c_j \in \mathbf{R}$, $j = 1, \cdots, 2n-1$, $c_{2n} > 0$, and the ultraviolet cutoff condition $\rho_x \in \mathcal{S}'_\mathrm{real}$ for each $\mathbf{x} \in \mathbf{R}^d$ is supposed.

By using the functional integral representations of e^{-tH_p} and e^{-tH_b}, the functional integral representation of e^{-tH_κ} is derived in the main theorem. Then, from the functional integral representation, it is seen that $e^{-tH\kappa}$ is positivity improving. Then, as a corollary of the main theorem, it is seen that the ground state of H_κ is unique if it exists.

For the spectral analysis for quantum particles systems coupled to Bose fields by the methods of stochastic analysis has been analyzed. For non-relativistic QED model, its functional representation is obtained in [8], and the case with spin is considered in [12]. The self-adjointness of the Hamiltonian is investigated in [11], the analysis of the bound state in [10] and that of the exponential decay in [7]. For spin-boson model and the Nelson model, the applications of their functional integral representation to spectral analysis are investigated in [2, 3, 9, 14, 18, 25, 26].

This paper is organized as follows. In section 2, the functional integral representation for the semi-relativistic particles is constructed. In section 3, we overview the Euclidean quantum field theory, and the functional integral representation for scalar Bose field is derived. In section 4, the interaction system is introduced. The the main theorem is stated, and its proof is given.

2 Relativistic Schrödinger Operator

According to [13], the functional integral representation of semigroup generated by the relativistic schrödinger operator is derived as follows. In this derivation, the Lévy subordinator plays an important role. A stochastic process $\{T_t\}_{t\geq 0}$ is called a Lévy subordinator if $\{T_t\}_{t\geq 0}$ is one dimensional Lévy process starting at zero and almost surely non-decreasing in $t \geq 0$.

The function $\Psi \in C^\infty((0,\infty))$ is called Bernstein function if $\Psi \geq 0$ and $(-1)^n \frac{d^n\Psi}{dx^n} \leq 0$ for all $n \in \mathbf{N}$. It is known in ([13] ; Proposition 2.5) that for a Bernstein function Ψ satisfying $\lim_{x\to+0} \Psi(x) = 0$, there exists an unique Lévy subordinator $\{T_t^\Psi\}_{t\geq 0}$ such that $\mathbb{E}[e^{-sT_t^\Psi}] = e^{-t\Psi(s)}$.

Let $M > 0$ be the fixed mass of the particle, and let us set $h_\mathrm{rel}(s) = \sqrt{s+M^2} - M$, $s > 0$. Since h_rel is a Bernstein function, it is seen that there exists a Lévy subordinator $\{T_t\}_{t\geq 0}$ on the probability space $(\Omega_\mathrm{rel}, \mathfrak{B}_\mathrm{rel}, P_\mathrm{rel})$ satisfying

$$\mathbb{E}_\mathrm{rel}[e^{-sT_t}] \;=\; e^{-th_\mathrm{rel}(s)}, \tag{3}$$

where $\mathbb{E}_{\rm rel}[X]=\int_{\Omega_{\rm rel}} X(\eta)dP_{\rm rel}(\eta)$.

Remark 2.1 *Let $Y_s=B_s+Ms$, $s>0$ where $\{B_s\}_{s>0}$ is one dimensional Brownian motion starting at zero. It is known that $\{T_t\}_{t\geq 0}$ is represented as the first hitting time process $T_t=\frac{1}{2}\inf\{s>0|Y_s=t\}$. (See [20]; Example 2.18).*

Let $\{\mathbf{B}_t\}_{t\geq 0}$ be d-dimensional Brownian motion starting $\mathbf{x}$ on the probability space $(\Omega_{\rm Br},\mathfrak{B}_{\rm Br},P^{\mathbf{x}}_{\rm Br})$. We introduce the probability space

$$(\Omega_{\rm p},\mathfrak{B}_{\rm p},P^{\mathbf{x}}_{\rm p})=(\Omega_{\rm rel}\times\Omega_{\rm Br},\overline{\mathfrak{B}_{\rm rel}\times\mathfrak{B}_{\rm Br}},\overline{P_{\rm rel}\times P^{\mathbf{x}}_{\rm Br}}),$$

and let us define a stochastic process $\{\mathbf{X}_t\}_{t\geq 0}$ on $(\Omega_{\rm p},\mathfrak{B}_{\rm p},P^{\mathbf{x}}_{\rm p})$ defined by

$$\mathbf{X}_t\left(\begin{bmatrix}\eta\\ \omega\end{bmatrix}\right)\;=\;\mathbf{B}_{T_t(\eta)}(\omega).$$

For $\psi\in L^2(\mathbf{R}^d_{\mathbf{x}})$, let us set

$$U^{\rm p}_t\psi(\mathbf{x})\;=\;\mathbb{E}^{\mathbf{x}}_{\rm p}[\psi(\mathbf{X}_t)e^{-\int_0^t V(\mathbf{X}_s)ds}],$$

where $\mathbb{E}^{\mathbf{x}}_{\rm p}[Z]=\int_{\Omega_{\rm p}} Z(\xi)dP^{\mathbf{x}}_{\rm p}(\xi)$. Here we assume the following condition.

(S.1) $V\in L^\infty(\mathbf{R}^d)$.

By Fourier transform and (3), it is seen that for rapidly decreasing function $\psi\in\mathcal{S}(\mathbf{R}^d)$,

$$(e^{-t(\sqrt{-\triangle+M^2}-M)}\psi)(\mathbf{x})=\int_{\mathbf{R}^d}e^{-th_{\rm rel}(\mathbf{k}^2)}\hat{\psi}(\mathbf{k})e^{i\mathbf{k}\cdot\mathbf{x}}d\mathbf{k}=\mathbb{E}_{\rm rel}[\mathbb{E}^{\mathbf{x}}_{\rm Br}[\psi(\mathbf{B}_{T_t})]]=\mathbb{E}^{\mathbf{x}}_{\rm p}[\psi(\mathbf{X}_t)].$$

Then from Trotter-Kato product formula, the functional integral representation for the semi-relativistic particle is obtained:

Proposition A ([13] ; Theorem 3.8)
Assume **(S.1)**. Then $(\phi,e^{-tH_{\rm p}}\psi)\;=(\phi,U^p_t\psi)$ for $\phi,\psi\in L^2(\mathbf{R}^d)$.

As a remark, in the proof of ([13]; Theorem 3.8) it is proven that a Feynman-Kac formula

$$e^{-t_1H_{\rm p}}g_1e^{-(t_2-t_1)H_{\rm p}}\cdots g_{n-1}e^{-(t_n-t_{n-1})H_{\rm p}}\psi(\mathbf{x})\;=\;\mathbb{E}^{\mathbf{x}}_{\rm p}[\prod_{j=1}^{n-1}g_j(\mathbf{X}_{t_j})\psi(\mathbf{X}_{t_n})e^{-\int_0^{t_n}V_{\rm p}(\mathbf{X}_s)ds}] \qquad (4)$$

holds where $g_j\in L^\infty(\mathbf{R}^d)$, $j=1,\cdots,n-1$ and $\psi\in L^2(\mathbf{R}^d)$.

3 Scalar Bose fields

3.1 Gaussian random process indexed by Hilbert space

In this subsection, basic properties for Gaussian random process are explained. To construct Bose fields, the following proposition is needed. (See, e.g.[1];Theorem 2.5, [20];Theorem 5.9)

Proposition B (Existence of Gaussian random process)
Let $\mathcal{K}$ be a separable and real Hilbert space. Then there exist a stochastic process $\{X_f\}_{f \in \mathcal{K}}$ indexed by $\mathcal{K}$ on a probability space $(Q_{\mathcal{K}}, \mathfrak{B}_{\mathcal{K}}, P_{\mathcal{K}})$ satisfying the following conditions.
(**G.1**) For all $f \in \mathcal{K}$, X_f is Gaussian random variable satisfying $\mathbb{E}[e^{-it\phi_f}] = e^{-\frac{\|f\|^2}{4}t^2}$.
(**G.2**) $X_{af+bg} = aX_f + bX_g$ for all $f, g \in \mathcal{K}$ and $a, b \in \mathbf{R}$.
(**G.3**) $\mathfrak{B}_{\mathcal{K}}$ is the minimal σ-field generated by $\{X_f\}_{f \in \mathcal{K}}$.

Remark 3.1 *The stochastic process $\{X_f\}_{f \in \mathcal{K}}$ satisfying **(G.1)-(G.3)** is called the Gaussian random process indexed by $\mathcal{K}$.*

Let $\{X_f\}_{f \in \mathcal{K}}$ be the Gaussian random process indexed by $\mathcal{K}$. Then it is seen from **(G.3)**, that $\mathcal{D}_{0,\mathcal{K}} = \left\{ F(X_{f_1}, \cdots, X_{f_n}) \,|\, F \in \mathcal{S}_{\mathrm{real}}(\mathbf{R}^d), f_j \in \mathcal{K}, j = 1, \cdots, n, n \in \mathbf{N} \right\}$ is dense in $L^2(Q_{\mathcal{K}})$. Let $L_n^2(Q_{\mathcal{K}})$ be the closure of the linear hull of the set $\{: \prod_{j=1}^n X_{f_j} : \,|\, f_j \in \mathcal{K},\, j = 1, \cdots, n\} \cup \{1\}$ where $: \prod_{j=1}^n X_{f_j} :$ denotes the wick product defined recursively by $: \prod_{j=1}^n X_{f_j} := X_{f_1} : \prod_{j=2}^n X_{f_j} : -\frac{1}{2} \sum_{j=2}^n (f_1, f_j) : \prod_{j \neq l} X_{f_l} :$ and $: X_f := X_f$. It is seen that $L_j^2(Q_{\mathcal{K}}) \perp L_l^2(Q_{\mathcal{K}})$ for $j \neq l$. It is known that the Winer-Ito-Segal decomposition $L^2(Q_{\mathcal{K}}) = \bigoplus_{n=0}^{\infty} L_n^2(Q_{\mathcal{K}})$ follows (See e.g. [1]; Lemma 2.13 , [20]; Lemma 5.4). Let S be a closed operator on $\mathcal{K}$. $\Gamma(S) = \oplus_{n=0}^{\infty} \Gamma^{(n)}(S)$ is called the second quantization of S defined by $\Gamma^{(n)}(S) X_{f_1} \cdots X_{f_n} \; = : X_{Sf_1} \cdots X_{Sf_n}$ for $f_j \in \mathcal{D}(S)$, $j = 1, \cdots, n$, $n \geq 0$. In addition, $d\Gamma(S) = \oplus_{n=0}^{\infty} d\Gamma^{(n)}(S)$ is called the differential second quantization of S defined by $d\Gamma^n(S) X_{f_1} \cdots X_{f_n} \; = \; \sum_{j=1}^n : X_{f_1} \cdots X_{Sf_j} \cdots X_{f_n} :$ where $f_j \in \mathcal{D}(S)$, $j = 1, \cdots, n$, $n \geq 0$.

3.2 Construction of a scalar Bose field

Let

$$\mathcal{K}_{\mathrm{b}}^0 \; = \; \left\{ f \in \mathcal{S}'_{\mathrm{real}}(\mathbf{R}^d) \,\middle|\, \int_{\mathbf{R}^d} \frac{|\hat{f}(\mathbf{k})|^2}{\omega_{\mathrm{b}}(\mathbf{k})} d\mathbf{k} \, \infty \right\},$$

where $\mathcal{S}'_{\mathrm{real}}(\mathbf{R}^d)$ is the space of real-valued tempered distributions, and set

$$(g, f)_{\mathcal{K}_{\mathrm{b}}} \; = \; \int_{\mathbf{R}^d} \frac{\overline{\hat{g}(\mathbf{k})}\, \hat{f}(\mathbf{k})}{\omega_{\mathrm{b}}(\mathbf{k})} d\mathbf{k}. \tag{5}$$

Here ω_{b} satisfies the following condition.

(B.1) $\omega_{\rm b}$ is continuous and non-negative.

As a remark, we consider a physical example of $\omega_{\rm b}$. Let $\omega_{\rm b}(\mathbf{k}) = \sqrt{\mathbf{k}^2 + m^2}$, where $m \geq 0$ denotes the mass of the field.

Let $\mathcal{K}_{\rm b} = \overline{\mathcal{K}_{\rm b}^0}^{\|\cdot\|_{\mathcal{K}_{\rm b}}}$. From Proposition B, there exists a Gaussian random process $\{\phi(f)\}_{f \in \mathcal{K}_{\rm b}}$ indexed by $\mathcal{K}_{\rm b}$ on a probability space $(Q_{\rm b}, \mathfrak{B}_{\rm b}, P_{\rm b})$. Let $\check{\omega}_{\rm b} = \omega_{\rm b}(-i\nabla)$. The free Hamiltonian of the bose field is given by

$$H_{\rm b} \ = \ d\Gamma(\check{\omega}_{\rm b}).$$

Then the triplet $(L^2(Q_{\mathcal{K}_{\rm b}}), H_{\rm b}, \{\phi(f)\}_{f \in \mathcal{K}_{\rm b}})$ of the scalar Bose field is constructed.

3.3 Functional integral representation for scalar Bose fields

In this subsection, we apply the Euclidean quantum filed theory. For the detail of this subject, refer to e.g. ([1]; Section 7) and ([20]; Section 5).

Let

$$\mathcal{K}_{\rm E}^0 \ = \ \left\{ f \in \mathcal{S}'_{\rm real}(\mathbf{R}^{1+d}) \,\middle|\, \int_{\mathbf{R}^{1+d}} \frac{|\hat{f}(k_0, \mathbf{k})|^2}{\omega_{\rm b}(\mathbf{k})^2 + k_0^2} dk_0 d\mathbf{k} \ < \ \infty \right\},$$

and

$$(g, f)_{\mathcal{K}_{\rm E}} \ = \ \int_{\mathbf{R}^d} \frac{\overline{\hat{g}(k_0, \mathbf{k})}\, \hat{f}(k_0, \mathbf{k})}{\omega_{\rm b}(\mathbf{k})^2 + k_0^2} dk_0 d\mathbf{k}. \tag{6}$$

Let $\mathcal{K}_{\rm E} = \overline{\mathcal{K}_{\rm E}^0}^{\|\cdot\|_{\mathcal{K}_{\rm E}}}$. From Proposition B, it is seen that there exists Gaussian random variables $\{\phi^{\rm E}(f)\}_{f \in \mathcal{K}_{\rm E}}$ indexed by $\mathcal{K}_{\rm E}$ on a probability space $(Q_{\rm E}, \mathfrak{B}_{\rm E}, P_{\rm E})$.

The relation between $\mathcal{K}_{\rm b}$ and $\mathcal{K}_{\rm E}$ is as follows. For the delta function $\delta_t \in \mathcal{S}'(\mathbf{R})$ with $< \delta_t, \phi >= \phi(t)$, it is seen that

$$(g, e^{-|t-s|\check{\omega}_{\rm b}} f)_{\mathcal{K}_{\rm b}} \ = \ (\delta_s \otimes g, \delta_t \otimes f)_{\mathcal{K}_{\rm E}}, \qquad s \neq t, \tag{7}$$

$$\|f\|_{\mathcal{K}_{\rm b}} \ = \ \|\delta_t \otimes f\|_{\mathcal{K}_{\rm E}}. \tag{8}$$

Then the isometric operator $j_t : \mathcal{K}_{\rm b} \to \mathcal{K}_{\rm E}$ is defined by $j_t f = \delta_t \otimes f$. Let $J_t = \Gamma(j_t)$. Then it is seen that $e^{-tH_{\rm b}} = J_0^* J_t$ and

$$(\Phi, e^{-tH_{\rm b}} \Psi)_{L^2(Q_{\rm b})} \ = \ \mathbb{E}_{\rm E}[(J_0 \Phi)^* (J_t \Psi)], \tag{9}$$

where $\mathbb{E}_{\rm E}(X) = \int_{Q_{\rm E}} X(\tilde{q}) dP_{\rm E}(\tilde{q})$. For $D \subset \mathbf{R}$, let us set $E_D = \Gamma(e_D)$ where e_D is the projection onto $\mathcal{K}_{\rm E}(D) = \{ f \in \mathcal{K}_{\rm E} | supp f \in D \times \mathbf{R}^d \}$. It is seen that E_D has the Markov property such that $E_{[a,b]} E_{\{c\}} E_{[d,e]} = E_{[a,b]} E_{[d,e]}$ for $a \leq b \leq c \leq d \leq e$. Let $E_s = J_s J_s^*$. Then it is known that $E_s = E_{\{s\}}$. It is seen that

$$J_s G(\phi(f)) J_s = E_s G(\phi^{\rm E}(\delta_s \otimes f)) E_s. \tag{10}$$

for $G \in L^\infty(\mathbf{R})$. Then by using Trotter-Kato product formula, the following proposition holds (Refer to e.g. [1]; Theorem 7.19).

Proposition C
Assume that V_b is continuous function on $\mathbf{R}^d$ with bounded from bellow. Then

$$(\Phi, e^{-t(H_\mathrm{b} \dot{+} V_\mathrm{b}(\phi(f)))}\Psi)_{L^2(Q_\mathrm{b})} \;=\; \mathbb{E}_\mathrm{E}\big[\overline{(J_0\Phi)}(J_t\Psi)e^{-\int_0^t V_\mathrm{b}(\phi^\mathrm{E}(\delta_s \otimes f))ds}\big].$$

4 Main Theorem and Proofs

4.1 Interacting system and main theorem

The interaction system between the semi-relativistic particle and a scalar Bose fields is defined as follows. The state space for the system is given by

$$\mathcal{H} \;=\; L^2(\mathbf{R}^d_\mathbf{x}) \otimes L^2(Q_\mathrm{b}).$$

$\mathcal{H}$ can be decomposed as $\mathcal{H} \simeq \int_{\mathbf{R}^d}^{\oplus} L^2(Q_\mathrm{b})d\mathbf{x}$ where $\int^{\oplus}$ denotes the fibre direct integral. The total Hamiltonian of the system is defined by form sum of the free Hamiltonian and interaction

$$H_\kappa \;=\; H_0 \,\dot{+}\, \kappa H_\mathrm{I}, \qquad \kappa \in \mathbf{R}, \tag{11}$$

where $H_0 = H_\mathrm{p} \otimes I + I \otimes H_\mathrm{b}$ and and H_I is given by

$$H_\mathrm{I} \;=\; \int_{\mathbf{R}^d}^{\oplus} P(\phi(\rho_\mathbf{x}))d\mathbf{x}$$

with $P(\lambda) = \sum_{j=1}^{2n} c_j\lambda^j$, $c_j \in \mathbf{R}$, $j = 1, \cdots, 2n-1$, $c_{2n} > 0$ and ρ_x satisfying the following conditions.

(A.1) For each $\mathbf{x} \in \mathbf{R}^d$, $f_\mathbf{x} \in \mathcal{K}_\mathrm{b}$ and $\sup_{\mathbf{x}\in\mathbf{R}^d} \|\rho_\mathbf{x}\|_{\mathcal{K}_\mathrm{b}} < \infty$.
(A.2) For each $t \in \mathbf{R}$, the map $\mathbf{R} \ni \mathbf{x} \mapsto \delta_t \otimes \rho_\mathbf{x} \in \mathcal{K}_\mathrm{E}$ is strongly continuous.

For a physical example of the interaction, let $\rho_\mathbf{x}(\mathbf{y}) = \rho(\mathbf{y}-\mathbf{x})$ for $\rho \in \mathcal{S}'_\mathrm{real}(\mathbf{R}^d)$. Then $\hat{\rho}_\mathbf{x}(\mathbf{k}) = \hat{\rho}(\mathbf{k})e^{i\mathbf{k}\cdot\mathbf{x}}$. Then we see that the conditions **(A.1)** and **(A.2)** are satisfied. The field operator $\phi(\rho_\mathbf{x})$ can be unrigorously represented as

$$\phi(\rho_\mathbf{x}) = \int_{\mathbf{R}^d} \frac{\hat{\rho}(\mathbf{k})}{\sqrt{2\omega(\mathbf{k})}}\left(a_\mathbf{k}e^{-i\mathbf{k}\cdot\mathbf{x}} + a_\mathbf{k}^\dagger e^{i\mathbf{k}\cdot\mathbf{x}}\right)d\mathbf{k}$$

where $a_\mathbf{k}$ and $a_\mathbf{k}^\dagger$ denote the kernel of an annihilation operator and creation operator, respectively.

We prepare for some notations. Let $\mathcal{D}_0 = C_0^\infty(\mathbf{R}^d)\hat{\otimes}\mathcal{D}_{0,\mathcal{K}_\mathrm{b}}$ where $\hat{\otimes}$ denotes the algebraic tensor product. For $\Psi \in \mathcal{H}$, we set $\Psi_{\mathbf{x}}(q) = \Psi(\mathbf{x}, q)$. Unless confusion arises, we identify $X \otimes I$ with X and $I \otimes Y$ with Y. Let

$$(\Omega_{\mathrm{p}\times\mathrm{E}}, \mathfrak{B}_{\mathrm{p}\times\mathrm{E}}, P^{\mathbf{x}}_{\mathrm{p}\times\mathrm{E}}) = (\Omega_{\mathrm{p}} \times Q_{\mathrm{E}}, \overline{\mathfrak{B}_{\mathrm{p}} \times \mathfrak{B}_{\mathrm{E}}}, \overline{P^{\mathbf{x}}_{\mathrm{p}} \times P_{\mathrm{E}}}),$$

and we use the notation $\mathbb{E}^{\mathbf{x}}_{\mathrm{p}\times\mathrm{E}}[Z] = \int_{\Omega_{\mathrm{p}\times\mathrm{E}}} Z(\zeta) dP^{\mathbf{x}}_{\mathrm{p}\times\mathrm{E}}(\zeta)$.

The main theorem in this paper is as follows.

Theorem 4.1 *Assume **(S.1)**, **(B.1)**, **(A.1)** and **(A.2)**. Then it follows that*

$$(\Phi, e^{-tH_\kappa}\Psi)_{\mathcal{H}} \ = \ \int_{\mathbf{R}^d} \mathbb{E}^{\mathbf{x}}_{p\times E}[\overline{(J_0\Phi_{\mathbf{X}_0})}(J_t\Psi_{\mathbf{X}_t})e^{-\int_0^t V(\mathbf{X}_s)ds} e^{-\kappa P(\phi^E(\int_0^t \delta_s \otimes \rho_{\mathbf{X}_s} ds)))}] d\mathbf{x}.$$

Now we consider an application of the above theorem. For a self-adjoint H with bounded from below, it is said that H has the ground state if the infimum of the spectrum of H is the eigenvalue. It is seen that $e^{-tH_{\mathrm{p}}}$ and $e^{-tH_{\mathrm{b}}}$ are positivity improving operators. Hence from the above functional integral representation, the next corollary immediately follows.

Corollary 4.2
*Assume **(S.1)**, **(B.1)**, **(A.1)** and **(A.2)**. Then if H_κ has the ground state, it is unique.*

4.2 Proof of Theorem 4.1

To prove the Theorem 4.1, we show the following proposition.

Proposition 4.3 *Let $G_j \in L^\infty(\mathbf{R}^d)$, $j = 1, \cdots n$. Then it follows that for $\Phi, \Psi \in \mathcal{D}_0$,*

$$\begin{aligned} &(\Phi, e^{-t_1 H_0} G_1(\phi(\rho_{\mathbf{x}})) e^{-(t_2 - t_1)H_0} G_2(\phi(\rho_{\mathbf{x}})) \cdots G_{n-1}(\phi(\rho_{\mathbf{x}})) e^{-(t_n - t_{n-1})H_0}\Psi) \\ &= \int_{\mathbf{R}^d} \mathbb{E}^{\mathbf{x}}_{p\times E} \left[\overline{(J_0 \Phi_{\mathbf{X}_{t_1}})} \left(\prod_{j=1}^{n-1} G_j(\phi(\delta_{t_j} \otimes \rho_{\mathbf{x}})) \right) (J_{t_n}\Psi_{\mathbf{X}_{t_n}}) e^{-\int_0^{t_n} V(\mathbf{X}_s)ds} \right] d\mathbf{x}. \end{aligned}$$

(Proof) By using $e^{-(t-s)H_{\mathrm{b}}} = J_s^* J_t$ for $t > s$ and (10), it is seen that

$$\begin{aligned} &(\Phi, e^{-t_1 H_0} G_1(\phi(\rho_{\mathbf{x}})) e^{-(t_2 - t_1)H_0} G_2(\phi(\rho_{\mathbf{x}})) \cdots G_{n-1}(\phi(\rho_{\mathbf{x}})) e^{-(t_n - t_{n-1})H_0}\Psi)_{\mathcal{H}} \\ &= \int_{\mathbf{R}^d} (\Phi_{\mathbf{x}}, e^{-t_1 H_0} G_1(\phi(\rho_{\mathbf{x}})) e^{-(t_2 - t_1)H_0} G_2(\phi(\rho_{\mathbf{x}})) \cdots G_{n-1}(\phi(\rho_{\mathbf{x}})) e^{-(t_n - t_{n-1})H_0}\Psi_{\mathbf{x}})_{L^2(Q_{\mathrm{b}})} d\mathbf{x} \\ &= \int_{\mathbf{R}^d} (\Phi_{\mathbf{x}}, e^{-t_1 H_{\mathrm{p}}} J_0^* \left(E_{t_1} G_1(\phi^{\mathrm{E}}(\delta_{t_1} \otimes \rho_{\mathbf{x}})) E_{t_1} \right) e^{-(t_2 - t_1)H_{\mathrm{p}}} \left(E_{t_2} G_2(\phi^{\mathrm{E}}(\delta_{t_2} \otimes \rho_{\mathbf{x}})) E_{t_2} \right) \times \\ &\qquad \cdots \times \left(E_{t_{n-1}} G_{n-1}(\phi^{\mathrm{E}}(\delta_{t_{n-1}} \otimes \rho_{\mathbf{x}})) E_{t_{n-1}} \right) J_{t_n} e^{-(t_n - t_{n-1})H_{\mathrm{p}}} \Psi_{\mathbf{x}})_{L^2(Q_{\mathrm{b}})} d\mathbf{x}. \end{aligned} \tag{12}$$

By using Markov property of E_{t_j} and the Feynman-Kac formula (4), we have

$$
\begin{aligned}
(12) = \int_{\mathbf{R}^d} \mathbb{E}_{\mathrm{E}}[(\overline{J_0\Phi_{\mathbf{x}}})e^{-t_1 H_{\mathrm{p}}} G_1(\phi^{\mathrm{E}}(\delta_{t_1}\otimes\rho_{\mathbf{x}}))e^{-(t_2-t_1)H_{\mathrm{p}}} G_2(\phi^{\mathrm{E}}(\delta_{t_2}\otimes\rho_{\mathbf{x}}))\times \\
\cdots\times G_{n-1}(\phi^{\mathrm{E}}(\delta_{t_{n-1}}\otimes\rho_{\mathbf{x}}))e^{-(t_n-t_{n-1})H_{\mathrm{p}}} J_{t_n}\Psi_{\mathbf{x}})]d\mathbf{x} \\
= \int_{\mathbf{R}^d} \mathbb{E}_{\mathrm{E}}[\mathbb{E}_{\mathrm{p}}^{\mathbf{x}}[(\overline{J_0\Phi_{\mathbf{x}}})\left(\prod_{j=1}^{n-1} G_j(\phi^{\mathrm{E}}(\delta_{t_j}\otimes\rho_{\mathbf{X}_{t_j}}))\right)(J_{t_n}\Psi_{\mathbf{X}_{t_n}})e^{-\int_0^{t_n} V(\mathbf{X}_s)ds}]]d\mathbf{x}.
\end{aligned}
$$

Thus the proof is obtained. ■

(Proof of Theorem 4.1)
Let $\Phi,\Psi\in\mathcal{D}_0$. By Proposition 4.3 and Trotter-Kato product formula we have

$$
\begin{aligned}
(\Phi, e^{-tH_\kappa}\Psi) &= \lim_{n\to\infty}(\Phi, (e^{-\frac{t}{n}H_0}e^{-\frac{t}{n}\kappa H_{\mathrm{I}}})^n\Psi) \\
&= \lim_{n\to\infty}\int_{\mathbf{R}^d} \mathbb{E}_{\mathrm{p}\times\mathrm{E}}^{\mathbf{x}}[(\overline{J_0\Phi_{\mathbf{X}_0}})(J_t\Psi_{\mathbf{X}_t})e^{-\kappa P(\phi^{\mathrm{E}}(\sum_{j=1}^{n}(\frac{t}{n})\delta_{tj/n}\otimes\rho_{\mathbf{X}_{tj/n}}))} e^{-\int_0^t V(\mathbf{X}_s)ds}]d\mathbf{x} \qquad (13)
\end{aligned}
$$

Here note that

$$
\begin{aligned}
\|\delta_{t+\varepsilon}\otimes\rho_{\mathbf{X}_{t+\varepsilon}} - \delta_t\otimes\rho_{\mathbf{X}_t}\|_{\mathcal{K}_{\mathrm{E}}} &\le \|(\delta_{t+\varepsilon}-\delta_t)\otimes\rho_{\mathbf{X}_{t+\varepsilon}}\|_{\mathcal{K}_{\mathrm{E}}} + \|\delta_t\otimes(\rho_{\mathbf{X}_{t+\varepsilon}}-\rho_{\mathbf{X}_t})\|_{\mathcal{K}_{\mathrm{E}}} \\
&\le \|(1-e^{\varepsilon\check{\omega}_{\mathrm{b}}})^{1/2}\rho_{\mathbf{X}_{t+\varepsilon}}\|_{\mathcal{K}_{\mathrm{b}}} + \|(\rho_{\mathbf{X}_{t+\varepsilon}}-\rho_{\mathbf{X}_t})\|_{\mathcal{K}_{\mathrm{b}}} \qquad (14)
\end{aligned}
$$

It is known that the map $s\to\mathbf{X}_s(\xi)$ is continuous for each $\xi\in\Omega_{\mathrm{p}}$ except finite points. Then from **(A.2)** and (14), the map $\mathbf{R}\ni t\mapsto\delta_t\otimes\rho_{\mathbf{X}_t}\in\mathcal{K}_{\mathrm{E}}$ is strongly continuous almost surely. Then from this continuity and (13), we have

$$
(\Phi, e^{-tH_\kappa}\Psi) = \int_{\mathbf{R}^d} \mathbb{E}_{\mathrm{p}\times\mathrm{E}}^{\mathbf{x}}[(\overline{J_0\Phi_{\mathbf{X}_0}})(J_t\Psi_{\mathbf{X}_t})e^{-\kappa P(\phi^{\mathrm{E}}(\int_0^t \delta_s\otimes\rho_{\mathbf{X}_s}ds)))} e^{-\int_0^t V(\mathbf{X}_s)ds}]d\mathbf{x}. \qquad (15)
$$

Since $\mathcal{D}_0$ is dense in $\mathcal{H}$, the proof is obtained. ■

Acknowledgments

It is a pleasure to thank Professor Fumio Hiroshima for his advice and comments.

References

[1] A. Arai, *Functional integral methods in quantum mathematical physics*, (in japanese) kyoritsu Publication, 2010.

[2] V. Betz, F. Hiroshima, J. Lőrinczi, R. A. Minlos and H. Sphon, Ground state properties of the Nelson Hamiltonian - A Gibbs measure-based approach, *Rev. Math. Phys.* **14** (2002) 173-198.

[3] V. Betz and F. Hiroshima, Gibbs measure with double stochastic integrals on path space, *Inf. Dim. Ana. Quantum Prob. Related Topics* **12** (2009) 135-152.

[4] R. Carmona, M. C. Masters and B. Simon, Relativistic Schrödinger operators: Asymptotic behavior of the eigenvalues, *J. Funct. Anal.* **91** (1990) 117-142.

[5] J. Glimm and A. Jaffe, *Quantum physics: A functional integral point of view*, Springer 1987.

[6] T. Hidaka, Existence of a ground state for the Nelson model with a singular perturbation, *J. Math. Phys.* **52** (2011) 022102.

[7] T. Hidaka and F. Hiroshima, Pauli-Fierz model with Kato-class potential and exponential decay, *Rev. Math. Phys.* **22** (2010) 1181-1208.

[8] F. Hiroshima, Functional integral representation of a model in quantum electrodynamics, *Rev. Math. Phys.* **9** (1997) 489-530.

[9] F. Hiroshima, Weak coupling limit removing an ultraviolet cut-off for a Hamiltonian of particles interacting with a scalar field *J. Math. Phys.* **40** (1999) 1215-1236.

[10] F. Hiroshima, Ground state of a model in nonrelativistic quantum electrodynamics II, *J. Math. Phys.* **41** (2000) 661-674.

[11] F. Hiroshima, Self-adjointness of the Pauli-Fierz Hamiltonian for arbitrary values of coupling constants, *Ann. Henri Poincaré* **3** (2002) 171-201.

[12] F. Hiroshima and J. Lőrinczi, Functional integral representation of the Pauli-Fierz model with spin $1/2$, *L. Funct. Anal.* **254** (2008), 2127-2185.

[13] F. Hiroshima, T. Ichinose and J. Lörinczi, Path integral representation for Schrödinger operators with Bernstein functions of Laplacian, (arxiv : 0906.0103.)

[14] F. Hiroshima, J. Lőrinczi and T. Takaesu, A probabilistic representation of the ground state expectation of fractional powers of the boson number operator in the Nelson model, arxiv: 1105.1850.

[15] T. Ichinose and H. Tamura, Imaginary-time path integral for lelativistic spinless particle in an electromagnetic field, *Commun. Math. Phys.* **105** (1986) 239-257.

[16] K. Kaleta and J. Lőrinczi, Analytic properties of fractional Schrödinger semigroups and Gibbs measures for symmetric stable processes, arxiv: 1011.2713.

[17] J. Lőrinczi, R. A. Minlos and H. Spohn, Infrared regular representation of the three dimensional massless Nelson model, *Lett. Math. Phys.* **59** (2002) 189-198.

[18] J. Lőrinczi, R. A. Minlos and H. Spohn, The infrared behaviour in Nelson's model of a quantum particle coupled to a massless scalar field, *Ann. Henri poincaré* **3** (2002) 1-28.

[19] J. Lőrinczi and J. Malecki, Spectral properties of the massless relativistic Harmonic oscillator, arxiv: 1006.3665.

[20] J. Lörinczi, F. Hiroshima and V. Betz, Feynman-Kac-type theorem and Gibbs Measure on path space - with applications to rigorous quantum field theory - 2011.

[21] M. Loss, T. Miyao and H. Spohn, Lowest energy states in nonrelativistic QED: Atoms and ions in motion, *J. Funct. Anal.* **243** (2007) 353-393.

[22] T. Miyao and H. Spohn. Spectral analysis of the semi-relativistic Pauli-Fierz Hamiltonian, *J. Funct. Anal.* **256** (2009) 2123-2156.

[23] O. Matte and E. Stockmeyer, Exponential localization of hydrogen-like atoms in relativistic quantum electrodynamics, *Commun. Math. Phys.* **295** (2010).

[24] B. Simon, *The* $P(\phi)_2$ *Euclidean (quantum) field theory*, Princeton University Press, 1974.

[25] H.Spohn, Ground state(s) of spin-boson Hamiltonian, *Commun. Math. Phys.* **123** (1989) , 277-304.

[26] H.Spohn, Ground state of quantum particle coupled to a scalar boson field, *Lett. Math. Phys.* **44** (1998), 9-16.

[27] T. Takaesu, Scaling limits for the system of semi-relativistic particles coupled to a scalar Bose field, *Lett. Math. Phys.* **97** (2011) 213-225.